

**WEST**

Generate Collection

L31: Entry 23 of 27

File: USPT

Jul 14, 1981

DOCUMENT-IDENTIFIER: US 4278594 A

TITLE: Process for separation and isolation of AHF, von Willebrand's ristocetin cofactor (VWF:RCF) and fibronectin from blood plasma

ABPL:

A process for the separation and isolation of AHF, fibronectin and von Willebrand's ristocetin cofactor from blood plasma by adding a sulfated mucopolysaccharide to blood plasma to a concentration of from about 0.05 to about 0.4 mg/ml of plasma, cooling the plasma to 0.degree. C.-15.degree. C., centrifuging the plasma to isolate a precipitate which forms and then removing the AHF-rich supernatant liquid from the precipitate.

ABPL:

The precipitate is washed in a slightly basic solution of dilute salts and then dissolved in a buffered solution of chaotropic salts of the Hofmeister series having an ionic strength from 0.01-1. Fibronectin is separated by chromatographic means from the dissolved precipitate which contains fibronectin and fibrinogen.

ABPL:

The dissolved precipitate may also be applied to a glass or plastic column containing a 4% agarose gel (E. G. Sepharose 4B or Biogel A-15m) in a 0.25 molar solution sodium phosphate titrated to pH 7.0 with any base, preferably TRIS. The von Willebrand's ristocetin cofactor elutes in the void volume of the column, whereas the fibronectin-rich fraction elutes later. Both fractions may be collected and concentrated.

ABPL:

in a preferred embodiment, 0.20 mg of sodium heparin per milliliter of human plasma is employed with the heparinized plasma solution cooled for 3 hours at a temperature of 2.degree. C.-4.degree. C.

PCPR:

This application is a continuation-in-part application of my co-pending application Ser. No. 49,965, now Pat. No. 4,210,580, filed June 19, 1979 entitled PROCESS FOR SEPARATION AND ISOLATION OF AHF AND FIBRONECTIN FROM BLOOD PLASMA.

BSPR:

This invention is concerned with a method for obtaining significant amounts of certain components of blood plasma.

BSPR:

The important components of blood plasma with which the process of this invention is primarily concerned are the anti-hemophilia factor (AHF), plasma fibronectin (cold-insoluble globulin-CIg) and von Willebrand's ristocetin cofactor. Components obtained according to the process of this invention are valuable in therapeutic applications. The AHF is known, of course, as the single significantly effective agent for use in treating hemophilia A.

BSPR:

It has been demonstrated in recent years that the fibronectin (CIg) component of human plasma is especially useful in speeding tissue recovery in trauma, particularly burns. The affinity of fibronectin to collagen and/or fibrin may be of significance in the mode of its action.

BSPR:

The infusion of plasma fractions enriched with fibronectin has been effective in reducing general sepsis from severe trauma and burns. It has been found that a reduction in the body's level of fibronectin occurs following trauma, and that the sooner fibronectin is restored to normal levels, the faster recovery takes place; presumably related to an increase in the removal of damaged tissues, reduction of infection, and general wound healing.

BSPR:

Of concern to the growing application of fibronectin therapy is the depletion of plasma available for the isolation of AHF. This has vastly increased the need for an effective and efficient large scale process for separating plasma fractions of predominating fibronectin from the fractions of predominantly AFH activity.

BSPR:

The present invention utilizes a number of individually known steps in combination with unique approaches to develop an overall process for the large scale isolation and purification of fibronectin (CIg), AHF, and von Willebrand's ristocetin cofactor from blood plasma.

BSPR:

The separation of such components by heparin precipitate has been carried out on only small scale levels of a milliliter of plasma up to one or two hundred milliliters of plasma. No satisfactory method has previously been developed for the processing of significant amounts of plasma such as over 0.5 liters and up to 1-5 liters.

BSPR:

It is well known that many unpredicted difficulties are usually encountered in attempting to scale up laboratory experimental procedures with plasma to even minimal commercial levels.

BSPR:

According to the present invention, there is provided a process for the separation of the fibronectin-rich (CIg) and von Willebrand's ristocetin cofactor fractions and the AHF-rich fraction of blood plasma by cryoprecipitation in the presence of a very specific concentration of certain sulfated mucopolysaccharides such as heparin, alginic acid sulfate, chitin sulfate, polymannuronic acid sulfate, chondroitin sulfate, and dextran sulfate and the subsequent purification of the fibronectin fraction by chromatographic separations utilizing a unique buffer system.

BSPR:

According to the process of the present invention, very significantly greater percentages of fibronectin, von Willebrand's ristocetin cofactor and AHF components are recovered compared to existing methods. In addition, the process of the present invention, while applicable for small scale plasma separations to which prior methods have generally been limited, is unexpectedly useful in separating AHF, von Willebrand's ristocetin cofactor and fibronectin fractions on a large scale, i.e., more than 0.5 liters of blood plasma, with amounts of 1-2 liters being accomplished with ease. Most significantly, the extremely high percentage recovery of AHF, von Willebrand's ristocetin cofactor and fibronectin is applicable even when using the process with the larger volumes of blood plasma.

BSPR:

Heparin is the preferred polysaccharide to use at concentrations of 0.05-0.4 mg/ml of plasma, preferably from 0.15 to 0.20 mg/ml, with the most preferred concentration being about 0.20 mg/ml.

BSPR:

The heparin is added to the plasma, which is then cooled to 0.degree. C.-15.degree. C. and preferably 2.degree.-4.degree. C. for at least one hour, preferably for at least three hours.

BSPR:

Approximately 80-90% of the fibronectin component and 95% or more of the von Willebrand's ristocetin cofactor component of the plasma is precipitated in this manner.

BSPR:

After washing, there is gently added to the precipitate a buffer solution of a chaotropic salt of the Hofmeister series having an ionic strength between 0.01-1, preferably 0.05-0.5 with 0.2 ionic strength as representative of the most used level. Chaotropic salts such as KSCN and KI buffered in the pH range of 6-8 are preferred, with a solution of 0.2 M KSCN, 0.1 M TRIS-Cl at pH 7.5 as most often used to dissolve the precipitate. The buffer is generally warmed to about 37.degree. C.

BSPR:

The dissolved fibronectin and fibrinogen protein complex is then separated chromatographically. The solution is applied to a DEAE cellulose (Whatman DE-23) column.

BSPR:

The fibrinogen is usually eluted with 0.05 M KSCN, 0.1 M TRIS-Cl at a pH of about 6-8, preferably about 7.5. When absorbance returns to a baseline level, a linear gradient of 0.05 M KSCN, 0.1 M TRIS-Cl at pH 7.5 (75-300 ml/chamber) is applied, and finally, a wash of 0.5 M KSCN, 0.1 M TRIS-Cl at a pH of about 7.5. Heparin separates from the fibronectin at the end of the gradient.

BSPR:

Also, the fibronectin can be eluted with a single buffer after the fibrinogen has been removed. Any buffer of at least 0.25 ionic strength and a pH range of 6-8 may be employed, such as 0.25 M NaH<sub>2</sub>PO<sub>4</sub>-TRIS pH 7.0 or 0.3 M KSCN, 0.1 M TRIS-Cl pH 7.5 and preferably 0.25 M Phosphate-TRIS pH 7.0 or 0.3 M KSCN, 0.1 M TRIS-Cl pH 7.5 to remove the fibronectin from the resin with a minimal amount of contaminating heparin. However, there is a 10-20% decrease in recovery of fibronectin at this phase. Approximately 45-50% recovery of fibronectin is usually obtained with either system.

BSPR:

As to the heparin supernatant fraction, each liter is mixed with an anion exchange resin, for example, 20-30 grams of DEAE cellulose with micro-crystalline binder (such as Heparasorb, General Diagnostics) for 10-15 minutes. The resin is then removed by constant flow centrifugation or macrofiltration.

BSPR:

The dissolved precipitate can also be applied alternatively to a glass or plastic column containing, for instance, a 4% agarose gel (i.e., Sepharose 4B or Biogel A-15m) in the foregoing buffer. The von Willebrand's ristocetin cofactor elutes in the void volume of the column first, and the fibronectin fraction is eluted later. Each fraction is collected and concentrated. In this alternative embodiment, from 95-100% of von Willebrand's ristocetin cofactor is obtained and from about 50-60% of the fibronectin.

BSPR:

In this specification and the claims, it should be understood that the term blood plasma includes blood plasma itself and cryoprecipitate which has been dissolved in dilute salt buffers.

DEPR:

1.5 liters of human blood plasma are warmed at 37.degree. C. for 30 minutes. 1.6 mls of a 200 mg/ml sodium heparin solution (sigma 167 u/mg) is mixed with the plasma and the plasma then cooled at 4.degree. C. for 18 hours (overnight) and a precipitate formed. The precipitate is collected by centrifugation at 4.degree. C. for 30 min. at 9,500 .times.g. The supernatant liquid is saved.

DEPR:

The diluted sample is applied to a 2.5.times.40 cm DEAE cellulose column pre-equilibrated with 0.05 M KSCN, 0.1 M TRIS-Cl pH 7.5 and the column washed with 0.05 M KSCN buffer until the absorbance is returned to baseline. A stepwise buffer, 0.25 M TRIS-PO<sub>4</sub>, pH 7.0 is used rather than a gradient to elute the

fibronectin. A total of 176.5 mg of fibronectin is eluted which represents about 40-45% recovery of total fibronectin, assuming a 250-300 mg/liter starting concentration.

## DEPR:

300 mls of human blood plasma are warmed to 37.degree. C. To the warm plasma, NaN.sub.3 is added to a dilution of 0.02%, and Trasylol to a dilution of 2 mg/ml. To this solution is added sodium heparin to obtain a final concentration of 0.2 mg/ml of heparin. The solution is stirred until the materials are completely dissolved. The solution is cooled in a water bath at 2.degree. C. for three hours and then centrifuged at 10,000 .times.g for 30 minutes at 2.degree. C. The supernatant liquid contains 90-95% of the AHF material.

## DEPR:

The dissolved precipitate is applied to a 1.5.times.30 cm DEAE cellulose column and the fibrinogen is eluted with a 0.05 M KI, 0.1 M TRIS-Cl pH 7.5 buffer. The fibronectin (CIg) is then eluted with a linear gradient of 0.05 M KI, 0.1 M TRIS-Cl pH 7.5 buffer. The fibronectin peak was pooled into three fractions adding up to 83 mg from a starting amount of 128 mg, yielding a 65% recovery.

## DEPR:

To 3 ml. of plasma, 0.2 mg of heparin is added for each milliliter and is cooled to 4.degree. C. for 3 hours. This causes the precipitation of 195.7 micrograms of fibronectin per ml. from the plasma which originally contained 242 micrograms of fibronectin per milliliter.

## DEPR:

To 3 ml. of plasma, 0.388 mg of heparin is added for each milliliter which results in a precipitation of 185.6 micrograms of fibronectin.

## DEPR:

To 3 ml. of plasma, 0.048 mg of heparin is added for each milliliter which results in a precipitation of 176.2 micrograms of fibronectin.

## CLPR:

1. A process for the separation and isolation of the AHF fraction, the fibronectin fraction and the von Willebrand's ristocetin cofactor fraction from blood plasma comprising:

## CLPR:

3. A process as claimed in claim 2, wherein said heparin is present in an amount of 0.15-0.25 mg/ml of plasma.

## CLPR:

4. A process as claimed in claim 2, wherein said heparin is present in an amount of 0.20 mg/ml of plasma.

## CLPR:

5. A process as claimed in any one of the preceding claims 1-4, wherein said blood plasma is cooled to 2.degree. C.-4.degree. C. for 30 minutes to 3 hours.

CLPR:

6. A process as claimed in any one of the preceding claims 1-4, wherein said chaotropic salt has an ionic strength of from 0.05-0.5.

CLPR:

7. A process for the separation and isolation of the AHF fraction, the fibronectin fraction and the von Willebrand's ristocetin cofactor fraction from blood plasma comprising:

CLPR:

8. A process for the separation and isolation of the AHF fraction, the fibronectin fraction and the von Willebrand's ristocetin cofactor fraction from blood plasma comprising:

CLPR:

9. A process for the separation and isolation of the AHF fraction from blood plasma comprising:

CLPR:

11. A process as claimed in claim 10 wherein said heparin is present in amount of 0.15-0.25 mg/ml of plasma.

CLPR:

12. A process as claimed in claim 10 wherein said heparin is present in the amount of 0.20 mg/ml of plasma.

CLPR:

13. A process as claimed in any one of the preceding claims 10-12 wherein said plasma is cooled to 2.degree. C.-4.degree. C. for 30 minutes to 3 hours.

CLPR:

14. A process for the separation and isolation of the AHF fraction, the fibronectin fraction and the von Willebrand's ristocetin cofactor fraction from blood plasma comprising:

CLPR:

16. A process as claimed in claim 15 wherein said heparin is present in an amount of 0.15-0.25 mg/ml of plasma.

CLPR:

17. A process as claimed in claim 15 wherein said heparin is present in an amount of 0.20 mg/ml of plasma.

CLPR:

18. A process as claimed in any one of the preceding claims 14-17 wherein said chaotropic salt has an ionic strength of from 0.05-0.5.

CLPV:

(a) adding a sulfated mucopolysaccharide to blood plasma to a concentration of 0.05-0.4 mg/ml of plasma;

CLPV:

(b) then cooling said plasma to 0.degree. C.-15.degree. C.;

CLPV:

(c) centrifuging the plasma to isolate a precipitate which forms;

CLPV:

(f) thereafter dissolving said precipitate in a buffered solution of chaotropic salts of the Hofmeister series having an ionic strength from 0.01-1;

CLPV:

(g) and thereafter separating said fibronectin from the dissolved precipitate which contains fibronectin, fibrinogen and von Willebrand's ristocetin cofactor by chromatographic means.

CLPV:

(a) adding a sulfated mucopolysaccharide to blood plasma to a concentration of 0.05-0.4 mg/ml of plasma;

CLPV:

(b) then cooling said plasma to 0.degree. C. to 15.degree. C.;

CLPV:

(c) centrifuging the plasma to isolate a precipitate which forms;

CLPV:

(f) thereafter dissolving said precipitate in a buffered solution of chaotropic salts of the Hofmeister series having an ionic strength from 0.01-1;

CLPV:

(g) and separating out said von Willebrand's ristocetin cofactor chromatographically utilizing an agarose gel whereby said von Willebrand's ristocetin cofactor elutes first and thereafter separating said fibronectin which elutes second.

CLPV:

(a) adding a sulfated mucopolysaccharide to blood plasma to a concentration of 0.05 -0.4 mg/ml of plasma;

CLPV:

(b) then cooling said plasma to 0.degree. C.-15.degree. C.;

CLPV:

(c) centrifuging the plasma to isolate a precipitate which forms;

CLPV:

(f) thereafter dissolving said precipitate in a buffered solution of chaotropic salts of the Hofmeister series having an ionic strength from 0.01-1;

CLPV:

(a) adding a sulfated mucopolysaccharide to blood plasma to a concentration of 0.05-0.4 mg/ml of plasma;

CLPV:

(b) then cooling said plasma to 0.degree. C.-15.degree. C.;

CLPV:

(c) centrifuging the plasma to isolate a precipitate which forms;

CLPV:

(a) adding a sulfated mucopolysaccharide to blood plasma to a concentration of 0.05-0.4 mg/ml of plasma;

CLPV:

(b) then cooling said plasma to 0.degree. C. to 15.degree. C.;

CLPV:

(c) centrifuging the plasma to isolate a precipitate which forms;

CLPV:

(f) thereafter dissolving said precipitate in a buffered solution of chaotropic salts of the Hofmeister series having an ionic strength of 0.01-1;